

Q-ADAPTED QUANTUM STOCHASTIC INTEGRALS AND DIFFERENTIALS IN FOCK SCALE

V. P. BELAVKIN AND M. F. BROWN

ABSTRACT. In this paper we first introduce the Fock-Guichardet formalism for the quantum stochastic (QS) integration, then the four fundamental processes of the dynamics are introduced in the canonical basis as the operator-valued measures, on a space-time σ -field $\mathfrak{F}_X$, of the QS integration. Then rigorous analysis of the QS integrals is carried out, and continuity of the QS derivative $\mathbf{D}$ is proved. Finally, Q-adapted dynamics is discussed, including Bosonic ($Q = \mathbf{I}$), Fermionic ($Q = -\mathbf{I}$), and monotone ($Q = \mathbf{O}$) quantum dynamics. These may be of particular interest to quantum field theory, quantum open systems, and quantum theory of stochastic processes.

1. INTRODUCTION

Non-commutative generalization of the Itô stochastic calculus, developed in [1], [2], [10], [14], [17] and [18] gave an adequate mathematical tool for studying the behavior of open quantum dynamical systems singularly interacting with a boson quantum-stochastic field. Quantum stochastic calculus also made it possible to solve an old problem of describing such systems with continuous observation and constructing a quantum filtration theory which would explain a continuous spontaneous collapse under the action of such observation [4], [5] and [8]. This gave examples of stochastic non-unitary, non-stationary, and even non-adapted evolution equations in a Hilbert space whose solution requires a proper definition of chronologically ordered quantum stochastic semigroups, and exponents of operators, by extending the notion of the multiple stochastic integral to non-commuting objects.

Here is the first part of an outline of the solution to this important problem by developing a *Q-adapted* form of the new quantum stochastic calculus constructed in [7] in a natural scale of Fock spaces. It is based on an explicit definition, introduced in [6], of the non-adapted quantum stochastic integral, as a non-commutative generalization of the Skorokhod integral [20] represented in the Fock space. The point derivative of the quantum stochastic calculus is discussed as an operator on the scaled Fock space, and consequentially the single integral operator-kernels are derived from the operator-kernels of the multiple stochastic integral. These quantum stochastic derivatives are then presented in an explicit Q-adapted form, and

Date: October 2010.

Key words and phrases. Quantum Stochastic Calculus, Non-adapted Stochastic Integrals, White Noise Analysis, Quantum Stochastic Processes.

we recover the Fermionic anti-commutation relation as well as the trivial Bosonic commutator of the usual adapted process.

The approach used here is similar in spirit to the kernel calculus of Maassen-Lindsay-Meyer [14], [17], however the difference is that all the main objects are constructed not in terms of kernels but in terms of operators represented in the Fock space. In addition we employ a much more general notion of multiple stochastic integral, non-adapted in general but focusing now on $\mathbb{Q}$ -adapted processes, which reduces to the notion of the kernel representation of an operator only in the case of a scalar (non-random) operator function under the integral. The possibility of defining a non-adapted single integral in terms of the kernel calculus was shown by Lindsay [15], but the notion of the multiple quantum-stochastic integral was introduced in [7].

2. RIGGED GUICHARDET-FOCK SPACE

Let $(\mathbb{X}, \lambda)$ be an essentially ordered space, that is, a measurable space $\mathbb{X}$ with a σ -finite measure $\lambda : \mathfrak{F}_{\mathbb{X}} \ni \Delta \mapsto \lambda(\Delta) \geq 0$ and an ordering relation $x \leq x'$ with the property that any n -tuple $(x_1, \dots, x_n) \in \mathbb{X}^n$ can be identified up to a permutation with a chain $\varkappa = \{x_1 < \dots < x_n\}$ modulo the product measure $\prod_{i=1}^n dx_i$ of $dx := \lambda(dx)$. In other words, we assume that the measurable ordering is almost total, that is, for any n the product measure of n -tuples $\mathbf{s} \in \mathbb{X}^n$ with components $(x_1, \dots, x_n)$ that are not comparable is zero. Hence, in particular, it follows that the measure λ on $\mathbb{X}$ is atomless and we may assume that this essentially total ordering on $\mathbb{X}$ is induced from the linear order in $\mathbb{R}_+$ by a measurable map $t : \mathbb{X} \rightarrow \mathbb{R}_+$ relatively to which λ is absolutely continuous with respect to the Lebesgue measure dt on $\mathbb{R}_+$ in the sense of admitting the disintegration

$$\langle f \circ t, 1_{\Delta} \rangle_{\lambda} := \int_{\Delta} f(t(x)) \lambda(dx) = \int_0^{\infty} f(t) \lambda_{\Delta}(t) dt \equiv \langle f, \lambda_{\Delta} \rangle.$$

Here 1_{Δ} is the indicator of any integrable subset $\Delta \subseteq \mathbb{X}$ and f is any essentially bounded function $f : \mathbb{R}_+ \rightarrow \mathbb{C}$ and $\Delta \mapsto \lambda_{\Delta}(t)$ is defined by duality as a positive measure on $\mathbb{X}$ for each $t \in \mathbb{R}_+$. In any case we will fix a map t that the above condition holds and $t(x) < t(x')$ if $x < x'$, interpreting $t(x)$ as the time at the point $x \in \mathbb{X}$. For example, $t(x) = t$ for $x = (\vec{x}, t)$ if $\mathbb{X} = \mathbb{R}^d \times \mathbb{R}_+$ is the $(d+1)$ -dimensional space-time with the casual ordering [3] and $dx = d\vec{x}dt$, where $d\vec{x}$ is the standard volume element on d -dimensional space $\mathbb{R}^d \ni \vec{x}$.

We shall identify the finite chains $\varkappa$ with increasingly indexed n -tuples $(x_1, \dots, x_n) \equiv \mathbf{s}$ with $x_i \in \mathbb{X}$, $x_1 < \dots < x_n$, denoting by $\mathcal{X} = \sum_{n=0}^{\infty} \mathcal{X}_n$ the set of all finite chains as the union of the sets

$$\mathcal{X}_n = \{\mathbf{s} \in \mathbb{X}^n : x_1 < \dots < x_n\}$$

with one-element $\mathcal{X}_0 = \{\emptyset\}$ containing the empty chain as a subset of $\mathbb{X}$: $\emptyset = \mathbb{X}^0$. We introduce a measure ‘element’ $d\varkappa = \prod_{x \in \varkappa} dx$ on $\mathcal{X}$ induced by the direct sum

$\oplus_{n=0}^{\infty} \lambda^{\otimes n}(\Delta_n)$, $\Delta_n \in \mathfrak{F}_{\mathbb{X}}^{\otimes n}$ of product measures $d\mathbf{s} = \prod_{i=1}^n dx_i$ on $\mathbb{X}^n$ with the unit mass $d\mathbf{x} = 1$ at the only atomic point $\mathbf{x} = \emptyset$.

Let $\{\mathfrak{H}_x : x \in \mathbb{X}\}$ be a family of Hilbert spaces $\mathfrak{H}_x$, let $\mathfrak{p}_0$ be an additive semigroup of nonnegative essentially measurable locally bounded functions $q : \mathbb{X} \rightarrow \mathbb{R}_+$ with zero included $0 \in \mathfrak{p}_0$, and let $\mathfrak{p}_1 = \{1 + q_0 : q_0 \in \mathfrak{p}_0\}$. For example, in the case $\mathbb{X} = \mathbb{R}^d \times \mathbb{R}_+$ by $\mathfrak{p}_1$ we mean the set of polynomials $q(x) = 1 + \sum_{k=0}^m c_k |\vec{x}|^k$ with respect to the modulus $|\vec{x}| = (\sum x_i^2)^{1/2}$ of a vector $\vec{x} \in \mathbb{R}^d$ with coefficients $c_k \geq 0$. We denote by $K_*(q)$ the Hilbert space of essentially measurable vector-functions $k : x \mapsto k(x) \in \mathfrak{H}_x$ which are square integrable with the weight $q \in \mathfrak{p}_1$:

$$\|k\|(q) = \left(\int \|k(x)\|_x^2 q(x) dx \right)^{1/2} < \infty.$$

With $q \geq 1$, any space $K_*(q)$ can be embedded into the Hilbert space $\mathcal{K}_* = K_*(1)$, and the intersection $\cap_{q \in \mathfrak{p}_1} K_*(q) \subseteq \mathcal{K}_*$ can be identified with the projective limit $K_+ = \lim_{q \rightarrow \infty} K_*(q)$. This follows from the facts that the function $\|k\|(q)$ is increasing: $q \leq p \Rightarrow \|k\|(q) \leq \|k\|(p)$, and so $K_*(p) \subseteq K_*(q)$, and that the set $\mathfrak{p}_1$ is directed in the sense that for any $q = 1 + r$ and $p = 1 + s$, $r, s \in \mathfrak{p}_0$, there is a function in $\mathfrak{p}_1$ majorizing q and p (we can take for example $q + p - 1 = 1 + r + s \in \mathfrak{p}_1$). In the case of polynomials $q \in \mathfrak{p}_1$ on $\mathbb{X} = \mathbb{R}^d \times \mathbb{R}_+$ the decreasing family $\{K_*(q)\}$, where $\mathfrak{H}_x = \mathbb{C}$, is identical with the integer Sobolev scale of vector fields $k : \mathbb{R}^d \rightarrow L^2(\mathbb{R}_+)$ with values $k(x)(t) = k(x, t)$ in the Hilbert space $L^2(\mathbb{R}_+)$ of square integrable functions on $\mathbb{R}_+$. If we replace $\mathbb{R}^d$ by $\mathbb{Z}^d$ and if we restrict ourselves to the positive part of the integer lattice $\mathbb{Z}^d$, then we obtain the Schwartz space in the form of vector fields $k \in K_+$.

The dual space K_*^- to K_+ is the space of generalized vector-functions $f(x)$ defining the continuous functionals

$$\langle f | k \rangle = \int \langle f(x) | k(x) \rangle dx, \quad k \in K_+.$$

It is the inductive limit $K_- = \lim_{q \rightarrow 0} K_*(q)$ in the opposite scale $\{K_*(q) : q \in \mathfrak{p}_-\}$, where $\mathfrak{p}_-$ is the set of functions $q : \mathbb{X} \rightarrow (0, 1]$ such that $1/q \in \mathfrak{p}_1$, which is the union $\cup_{q \in \mathfrak{p}_-} K_*(q)$ of the inductive family of Hilbert spaces $K_*(q)$, $q \in \mathfrak{p}_-$, with the norms $\|k\|(q)$, containing as the minimal the space $\mathcal{K}_*$. Thus we obtain the Gel'fand chain

$$K_+ \subseteq K_*(q_+) \subseteq \mathcal{K}_* \subseteq K_*(q_-) \subseteq K_-$$

in the extended scale $\{K_*(q) : q \in \mathfrak{p}\}$, where $\mathfrak{p} = \mathfrak{p}_- \cup \mathfrak{p}_1$, with $q_+ \in \mathfrak{p}_1$, $q_- \in \mathfrak{p}_-$. The dual space $K_+^* = K_-^-$ is the space of the continuous linear functionals on K_+ containing the Hilbert space $\mathcal{K}$ called the *rigged space* with respect to the dense subspace $K^+ = K_-^*$ of $\mathcal{K}$ equipped with the projective convergence in the scale $\|k^*\|(q) = \|k\|(q)$ for $q \in \mathfrak{p}_1$.

We can similarly define a Fock-Gel'fand triple $(F_+, \mathcal{F}_*, F_-)$ with

$$F_+ = \cap_{q \in \mathfrak{p}_1} F_*(q), \quad \mathcal{F}_* = F_*(1), \quad F_- = \cup_{q \in \mathfrak{p}_-} F_*(q),$$

for the Hilbert scale $\{F_\star(q) : q \in \mathfrak{p}\}$ of the symmetric Fock spaces $F_\star(q) = \oplus_{n=0}^\infty K_\star^{(n)}(q)$ over $K_\star(q)$, where $K_\star^{(0)}(q) = \mathbb{C}$, $K_\star^{(1)}(q) = K_\star(q)$, and each $K_\star^{(n)}(q)$ for $n > 1$ is given by the product-weight $q_n(x_1, \dots, x_n) = \prod_{i=1}^n q(x_i)$ on $\mathbb{X}^n$. We shall consider the Guichardet [11] representation of the symmetric tensor-functions $\psi_n \in K_\star^{(n)}(q)$ regarding them as the restrictions $\psi|_{\mathcal{X}_n}$ of the functions $\psi : \varkappa \mapsto \psi(\varkappa) \in K_\star^\otimes(\varkappa)$ with sections in the Hilbert products $K_\star^\otimes(\varkappa) = \otimes_{x \in \varkappa} \mathfrak{k}_x$, square integrable with the product weight $q(\varkappa) = \prod_{x \in \varkappa} q(x)$:

$$\|\psi\|(q) = \left(\int \|\psi(\varkappa)\|^2 q(\varkappa) d\varkappa \right)^{1/2} < \infty.$$

The integral here is over all chains $\varkappa \in \mathcal{X}$ and defines the pairing on F_+ by

$$\langle \psi | \psi \rangle = \int \langle \psi(\varkappa) | \psi(\varkappa) \rangle d\varkappa, \quad \psi \in F_+.$$

In more detail we can write this in the form

$$\int \|\psi(\varkappa)\|^2 q(\varkappa) d\varkappa = \sum_{n=0}^\infty \int_{0 \leq t_1 < \dots < t_n < \infty} \dots \int \|\psi(x_1, \dots, x_n)\|^2 \prod_{i=1}^n q(x_i) dx_i,$$

where the n -fold integrals for $\psi_n \in K_\star^{(n)}$ are taken over simplex domains $\mathcal{X}_n = \{\mathbf{s} \in \mathbb{X}^n : t(x_1) < \dots < t(x_n)\}$.

One can easily establish an isomorphism between the space $F_\star(q)$ and the symmetric (or antisymmetric) Fock space over $K_\star(q)$ with a nonatomic measure dx in $\mathbb{X}$. It is defined by the isometry

$$\|\psi\|(q) = \left(\sum_{n=0}^\infty \frac{1}{n!} \int \dots \int \|\psi(x_1, \dots, x_n)\|^2 \prod_{i=1}^n q(x_i) dx_i \right)^{1/2},$$

where the functions $\psi(x_1, \dots, x_n)$ can be extended to the whole of $\mathbb{X}^n$ in a symmetric (or antisymmetric) way uniquely up to the measure zero due to nonatomicity of dx on $\mathbb{X}$.

3. EXPLICIT DEFINITION OF QS INTEGRALS

Let $\mathfrak{h}$ be a Hilbert space called the initial space for the Hilbert products $\mathcal{H}_\star = \mathfrak{h} \otimes \mathcal{K}_\star$ and $\mathcal{G}_\star = \mathfrak{h} \otimes \mathcal{F}_\star$. We consider the Hilbert scale $G_\star(q) = \mathfrak{h} \otimes F_\star(q)$, $q \in \mathfrak{p}$ of complete tensor products of $\mathfrak{h}$ and the Fock spaces over $K_\star(q)$, and we put

$$G_+ = \cap G_\star(q), \quad G_- = \cup G_\star(q)$$

which constitute the Gel'fand triple $G_+ \subseteq \mathcal{G}_\star \subseteq G_-$ dual to $G^+ \subseteq \mathcal{G} \subseteq G^-$ of the Hermitian adjoint bra-spaces $G^+ = G_+^*$, $\mathcal{G} = \mathcal{G}_\star^*$, $G^- = G_-^*$.

Let $(D_\nu^\mu)_{\nu=\circ, +}^{\mu=-, \circ}$ be a quadruple of functions D_ν^μ on $\mathbb{X}$ with kernel values $D_\nu^\mu(x) : G_+ \rightarrow G_-$ for $\mathfrak{k}_x = \mathbb{C}$, or, if $\mathfrak{k}_x \neq \mathbb{C}$, as continuous operators

$$(3.1) \quad \begin{aligned} D_+^-(x) : G_+ &\rightarrow G_-, & D_\circ^\circ(x) : \mathfrak{k}_x \otimes G_+ &\rightarrow \mathfrak{k}_x \otimes G_-, \\ D_+^\circ(x) : \mathfrak{k}_x \otimes G_+ &\rightarrow G_-, & D_\circ^-(x) : \mathfrak{k}_x \otimes G_+ &\rightarrow G_-. \end{aligned}$$

The continuity means that there is a $q \in \mathfrak{p}_1$ such that these operators are bounded from $G_\star(q) \supseteq G_+$ to $G(q)^\star \subseteq G_-$, where $G(q)^\star = G_\star(q^{-1})$. We assume that $D_+^-(x)$ is locally integrable in the sense that

$$\exists q \in \mathfrak{p}_1 : \|D_+^-\|_{q,t}^{(1)} = \int_{\mathbb{X}^t} \|D_+^-(x)\|_q dx < \infty, \quad \forall t < \infty,$$

where $\mathbb{X}^t = \{x \in \mathbb{X} : t(x) < t\}$, and

$$\|D\|_q = \sup_{\chi \in G_\star(q)} \{\|D\chi\|(q^{-1})/\|\chi\|(q)\}$$

is the norm of the continuous operator $D : G_\star(q) \rightarrow G_\star(q^{-1})$ which defines a bounded Hermitian form $D(\chi; \chi) := \langle \chi | D\chi \rangle$ on $G_\star(q)$. We also assume that $D_\circ^\circ(x)$ is locally bounded with respect to a strictly positive function s of x such that $1/s \in \mathfrak{p}_0$ in the sense that

$$\exists q \in \mathfrak{p}_1 : \|D_\circ^\circ\|_{q,t}^{(\infty)}(s) = \text{ess sup}_{x \in \mathbb{X}^t} \{s(x)\|D_\circ^\circ(x)\|_q\} < \infty, \quad \forall t < \infty;$$

here $\|D\|_q$ is the norm of the operator $D : \mathfrak{k}_x \otimes G_\star(q) \rightarrow \mathfrak{k}_x \otimes G_\star(q^{-1})$. Finally, we assume that $D_+^\circ(x)$ and $D_\circ^-(x)$ are locally square integrable with strictly positive function $r(x)$ such that $1/r \in \mathfrak{p}_0$, in the sense that

$$\exists q \in \mathfrak{p}_1 : \|D_+^\circ\|_{q,t}^{(2)}(r) < \infty, \quad \|D_\circ^-\|_{q,t}^{(2)}(r) < \infty, \quad \forall t < \infty,$$

where $\|D\|_{q,t}^{(2)}(r) = (\int_{\mathbb{X}^t} \|D(x)\|_q^2 r(x) dx)^{1/2}$ and $\|D\|_q$ are the respective norms of the operators

$$D_+^\circ(x) : G_\star(q) \rightarrow \mathfrak{k}_x \otimes G_\star(q^{-1}), \quad D_\circ^-(x) : \mathfrak{k}_x \otimes G_\star(q) \rightarrow G_\star(q^{-1}).$$

Then for any $t \in \mathbb{R}_+$ we can define a *generalized quantum stochastic (QS) integral*

$$(3.2) \quad \mathbf{i}_0^t(\mathbf{D}) = \int_{\mathbb{X}^t} \Lambda(\mathbf{D}, dx), \quad \Lambda(\mathbf{D}, \Delta) = \sum_{\mu, \nu} \Lambda_\nu^\mu(D_\nu^\mu, \Delta)$$

introduced in [18] as the sum of four continuous operators $\Lambda_\mu^\nu(D_\nu^\mu) : G_+ \rightarrow G_-$ described as operator measures on $\mathfrak{F}_\mathbb{X} \ni \Delta$ for $\Delta = \mathbb{X}^t$ with values

$$\begin{aligned} [\Lambda_-^+(D_+^-, \Delta)\chi](\vartheta) &= \int_{\Delta} [D_+^-(x)\chi](\vartheta) dx && \text{(preservation),} \\ [\Lambda_\circ^+(D_+^\circ, \Delta)\chi](\vartheta) &= \sum_{x \in \Delta \cap \vartheta} [D_+^\circ(x)\chi](\vartheta \setminus x) && \text{(creation),} \\ [\Lambda_-^\circ(D_\circ^-, \Delta)\chi](\vartheta) &= \int_{\Delta} [D_\circ^-(x)\dot{\chi}(x)](\vartheta) dx && \text{(annihilation),} \\ (3.3) \quad [\Lambda_\circ^\circ(D_\circ^\circ, \Delta)\chi](\vartheta) &= \sum_{x \in \Delta \cap \vartheta} [D_\circ^\circ(x)\dot{\chi}(x)](\vartheta \setminus x) && \text{(exchange).} \end{aligned}$$

Here $\chi \in G_+$, $\vartheta \setminus x = \{x' \in \vartheta : x' \neq x\}$ denotes the chain $\vartheta \in \mathcal{X}$ from which the point $x \in \vartheta$ has been eliminated, and $\dot{\chi}(x) \in \mathfrak{k}_x \otimes G_+$ is the *single point split* $\dot{\chi}(x) = \nabla_x \chi$,

or point derivative, defined for each $\chi \in G_+$ almost everywhere (namely, for $\vartheta \in \mathcal{X}$: $x \notin \vartheta$) as the function

$$[\nabla_x \chi](\vartheta) = \chi(\vartheta \sqcup x) \equiv \dot{\chi}(x, \vartheta),$$

where the operation $\varkappa \sqcup x$ denotes the disjoint union $\vartheta = \varkappa \cup x$, $\varkappa \cap x = \emptyset$ of chains $\varkappa \in \mathcal{X}$ and $x \in \mathbb{X} \setminus \varkappa$ with pairwise comparable elements. Note that the point splitter ∇ represents the Malliavin derivative [16] densely defined in Fock-Guichardet space as the bosonic annihilation operator $b(x) : G_+ \rightarrow \mathfrak{k}_x \otimes G_+$ by $[b(x)\chi](\vartheta) = \dot{\chi}(x, \vartheta)$ where one can take $\dot{\chi}(x, \vartheta) = 0$ if $x \in \vartheta$, and its inverse operator at x is $[\nabla_x^* \psi](\vartheta) = \psi(x, \vartheta \setminus x)$ with $[\nabla_x^* \psi](\vartheta) = 0$ if $x \notin \vartheta$ defines in this representation the Skorokhod non-adapted integral as the creation point integral

$$[\nabla^* \psi](\vartheta) = \sum_{x \in \vartheta} \psi(x, \vartheta \setminus x)$$

for any $\psi \in K_- \otimes G_-$. The continuity of this derivative as the projective limit map $G_+ \rightarrow K_+ \otimes G_+$ and the point integral as the adjoint map $K_- \otimes G_- \rightarrow G_-$ will simply follow from the isometricity of the multiple point splitter and co-isometricity of the adjoint multiple point integral as defined below, originally established in [7].

4. SPLIT OPERATOR AND ITS PROPERTIES

As it is proved below, we can consider the multiple bosonic annihilation operators $b^\otimes(v) : \chi \mapsto \dot{\chi}(v)$ eliminating several points v in $\mathcal{X}_n$, with $[b^\otimes(v)\chi](\vartheta) = 0$ if $v \subseteq \vartheta$, as partial isometries on the projective limit G_+ into $G_+^{(n)} = K_+^{\otimes n} \otimes G_+$. They are described for each $v = \{x_1, \dots, x_n\}$ in terms of the *n-point split*

$$(4.1) \quad \left[\Delta_v^{(n)} \chi \right](\vartheta) := \chi(\vartheta \sqcup v) \equiv \chi^{(n)}(v, \vartheta), \quad v \in \mathcal{X}_n,$$

where we denoted $\dot{\chi}(v) = \chi^{(n)}(v)$ for $n = |v|$. It is defined almost everywhere ($\vartheta \cap v = \emptyset$) on $\vartheta \in \mathcal{X}$ as the n -th order $\Delta_{v_n} = \otimes_{x \in v_n} \nabla_x$ point (or Malliavin) derivative [16] such that $\Delta^{(0)} = \text{I}$ and $\Delta^{(1)} = \nabla$. These n -tuple annihilations, densely defined as operators from G_+ into $\mathfrak{k}^\otimes(v) \otimes G_+$, are not continuous for each $v \in \mathcal{X}_n$ (except $v = \emptyset$ corresponding to $n = 0$ for which $b^\otimes(\emptyset) = \text{I}$), but they define projective-continuous linear maps into the space $G_+^{(n)}$ of functions $v \mapsto \psi(v)$ on $\mathcal{X}_n \subset \mathcal{X}$ for each $n \in \mathbb{N}$, and therefore have the adjoints $G_+^{(n)} \rightarrow G_+$. This follows from the projective contractivity of the maps $\Delta^{(n)} : G_+ \rightarrow G_+^{(n)}$, and their adjoints, such that each function $\chi^{(n)} = \Delta^{(n)} \chi$ is square-integrable on $\mathcal{X}_n$ with any $q_0 \in \mathfrak{p}_0$ being a component of the isometric operator

$$\Delta \chi = \int_{\mathcal{X}}^{\oplus} \Delta_v \chi dv = \oplus_{n=0}^{\infty} \Delta^{(n)} \chi \equiv \dot{\chi}.$$

The projective isometricity of the linear operator $\Delta = \int_{\mathcal{X}}^{\oplus} \Delta_v dv$, called the *multiple point splitter*,

$$\Delta : G_\star(q_0 + q_1) \rightarrow F_\star(q_0) \otimes G_\star(q_1)$$

is established in the following lemma.

Lemma. *The linear map $\Delta : \chi \mapsto [\Delta^{(n)}\chi]$ defined as $\Delta\chi = \oplus_{n=0}^{\infty} \Delta^{(n)}\chi$ in (4.1) for all $v \in \mathcal{X}$ is a projective isometry on Hilbert scale $\{G_{\star}(q) : q \in \mathfrak{p}\}$ into the scale $\{F_{\star}(q_0) \otimes G_{\star}(q_1) : q_0 \in \mathfrak{p}_0, q_1 \in \mathfrak{p}_1\}$ such that*

$$\|\Delta\chi\|(q_0, q_1) = \|\chi\|(q_0 + q_1).$$

The adjoint co-isometric operator $\langle \Delta^*\psi | \chi \rangle = \langle \psi | \Delta\chi \rangle$, defined on $\psi \in F_{\star}(q_0) \otimes G_{\star}(q_1)$ as the multiple point integral $\Delta^* = \sum \Delta_n^*$, is a contraction from $F_{\star}(q_0^{-1}) \otimes G_{\star}(q_1^{-1})$ into any $G_{\star}(q^{-1})$ with $q \geq q_0 + q_1$ such that $(\Delta_n^*)^* = \Delta^{(n)}$. In particular,

$$(4.2) \quad [\Delta_n^*\psi](\vartheta) = \sum_{\mathcal{X}_n \ni v \subseteq \vartheta} \psi(v, \vartheta \setminus v), \quad \vartheta \in \mathcal{X}$$

defines for $\psi(v, \varkappa) = \psi_n(v) \otimes \chi(\varkappa)$ the n -th order Skorokhod integral

$$[S_n(\psi_n)\chi](\vartheta) = \sum_{\mathcal{X}_n \ni v \subseteq \vartheta} \psi_n(v) \otimes \chi(\vartheta \setminus v) = [\Delta_n^*(\psi_n \otimes \chi)](\vartheta)$$

of $\psi_n \in K_{\star}^{(n)}(q_0^{-1})$ on $\chi \in G_{\star}(q_1^{-1})$.

Proof. We first of all establish the principal formula of the multiple integration

$$(4.3) \quad \int \sum_{v \subseteq \vartheta} f(v, \vartheta \setminus v) d\vartheta = \iint f(v, \varkappa) dv d\varkappa, \quad \forall f \in L^1(\mathcal{X} \times \mathcal{X}),$$

which will allow us to define the adjoint operator Δ^* . Let $f(v, \varkappa) = g(v)h(\varkappa)$ be the product of integrable complex functions on $\mathcal{X}$ of the form $g(v) = \prod_{x \in v} g(x)$, $h(\varkappa) = \prod_{x \in \varkappa} h(x)$ for any $v, \varkappa \in \mathcal{X}$. Employing the binomial formula

$$\sum_{v \subseteq \vartheta} g(v)h(\vartheta \setminus v) = \sum_{v \sqcup \varkappa = \vartheta} \prod_{x \in v} g(x) \prod_{x \in \varkappa} h(x) = \prod_{x \in \vartheta} (g(x) + h(x)),$$

and also the equality $\int f(v) dv = \exp\{\int f(x) dx\}$ for $f(v) = \prod_{x \in v} f(x)$, we obtain the formula

$$\int \sum_{v \subseteq \vartheta} g(v)h(\vartheta \setminus v) d\vartheta = \exp\left\{\int (g(x) + h(x)) dx\right\} = \iint g(v)h(\varkappa) dv d\varkappa,$$

which proves (4.3) on a set of product-functions f dense in $L^1(\mathcal{X} \times \mathcal{X})$.

Applying this formula to the scalar product $\langle \psi(v, \varkappa) | \psi(v, \varkappa) \rangle \in L^1(\mathcal{X} \times \mathcal{X})$, we obtain

$$\int \sum_{v \subseteq \vartheta} \langle \psi(v, \vartheta \setminus v) | \chi(\vartheta) \rangle d\vartheta = \iint \langle \psi(v, \varkappa) | \chi(v \sqcup \varkappa) \rangle dv d\varkappa,$$

that is, $\langle \Delta^*\psi | \chi \rangle = \langle \psi | \Delta\chi \rangle$, where $[\Delta\chi](v, \varkappa) = \chi(\varkappa \sqcup v) \equiv \hat{\chi}(v, \varkappa)$. Choosing arbitrary $\psi \in F_{\star}(q_0^{-1}) \otimes G_{\star}(q_1^{-1})$, we find that the annihilation operators $b(v)\chi = [\Delta_v\chi]$ define the isometry $\Delta : G_{\star}(q_0 + q_1) \rightarrow F_{\star}(q_0) \otimes G_{\star}(q_1)$ with the operator Δ^*

defined as co-isometry $F_*(q_0^{-1}) \otimes G_*(q_1^{-1}) \rightarrow G_*(q^{-1})$ for $q = q_0 + q_1$ with respect to the standard pairing of dual spaces $G_*(q)$ and $G_*(q^{-1})$:

$$\begin{aligned} \|\dot{\chi}\|^2(q_0, q_1) &= \iint \|\dot{\chi}(v, \varkappa)\|^2 q_0(v) q_1(\varkappa) dv d\varkappa \\ &= \int \sum_{v \subseteq \vartheta} \|\chi(\vartheta)\|^2 q_0(v) q_1(\vartheta \setminus v) d\vartheta = \int \|\chi(\vartheta)\|^2 \sum_{v \sqcup \varkappa = \vartheta} q_0(v) q_1(\varkappa) d\vartheta \\ &= \int \|\chi(\vartheta)\|^2 (q_0 + q_1)(\vartheta) d\vartheta \equiv \|\chi\|^2(q_0 + q_1) \leq \|\chi\|^2(q) \quad \forall q \geq q_0 + q_1. \end{aligned}$$

Hence it follows that Δ is projective continuous operator from G_+ to $F_+ \otimes G_+$, where $F_+ = \bigcap_{q \in \mathfrak{p}_0} F_*(q)$, and in particular so is the one-point split $\chi(x, \varkappa) = \chi(x \sqcup \varkappa) \equiv \dot{\chi}(x, \varkappa)$ from G_+ to $K_+ \otimes G_+$, as a contracting map $G_*(q_0 + q_1) \rightarrow F_*(q_0) \otimes G_*(q_1)$ for all $q_0 \in \mathfrak{p}_0, q_1 \in \mathfrak{p}$. The lemma is proved. $\square$

Remark. *Because the explicit form of both the creation and annihilation operators' norms may not be obvious we shall review them here for the reader's familiarization.*

$$\|\Delta\|_{q_0+q_1} = \sup_{\chi} \frac{\|\Delta\chi\|(q_0, q_1)}{\|\chi\|(q_0 + q_1)} = \sup_{\chi, \psi} \frac{|\langle \psi | \Delta\chi \rangle|}{\|\psi\|(q_0^{-1}, q_1^{-1}) \|\chi\|(q_0 + q_1)}$$

and

$$\|\Delta^*\|^{q_0, q_1} = \sup_{\psi} \frac{\|\Delta^*\psi\|((q_0 + q_1)^{-1})}{\|\psi\|(q_0^{-1}, q_1^{-1})} = \sup_{\chi, \psi} \frac{|\langle \Delta^*\psi | \chi \rangle|}{\|\psi\|(q_0^{-1}, q_1^{-1}) \|\chi\|(q_0 + q_1)}$$

where $\psi \in F_*(q_0^{-1}) \otimes G_*(q_1^{-1})$ and $\chi \in G_*(q_0 + q_1)$, indeed $\Delta^* : F_*(q_0^{-1}) \otimes G_*(q_1^{-1}) \rightarrow G_*((q_0 + q_1)^{-1})$ and $\Delta : G_*(q_0 + q_1) \rightarrow F_*(q_0) \otimes G_*(q_1)$. By virtue of the fact that $\langle \psi | \Delta\chi \rangle = \langle \Delta^*\psi | \chi \rangle$ these two norms are equal, and they are equal to 1.

5. MULTIPLE QS INTEGRALS AND THEIR CONTINUITY

We are now ready to prove the *inductive continuity* of the integral (3.2) with respect to $\mathbf{D} = [\mathbf{D}_\nu^\mu]$ by showing the inequality

$$\|(\mathbf{i}_0^t(\mathbf{D})\chi)\|(p^{-1}) \leq \|\mathbf{D}\|_{q,t}^s(r) \|\chi\|(p), \quad \forall p \geq r^{-1} + q + s^{-1},$$

where $\|\mathbf{D}\|_{q,t}^s(r) = \|\mathbf{D}_+^-\|_{q,t}^{(1)} + \|\mathbf{D}_+^\circ\|_{q,t}^{(2)}(r) + \|\mathbf{D}_\circ^-\|_{q,t}^{(2)}(r) + \|\mathbf{D}_\circ^\circ\|_{q,t}^{(\infty)}(s)$. We will establish this inequality as the single-integral case of the corresponding inequality for the *generalized multiple QS integral* [7]

$$(5.1) \quad [\mathbf{i}_0^t(\mathbf{M})\chi](\vartheta) = \sum_{v_\circ^- \sqcup v_+^\circ \subseteq \vartheta^t} \int_{\mathcal{X}^t} \int_{\mathcal{X}^t} [\mathbf{M}(\mathbf{v}) \dot{\chi}(v_\circ^- \sqcup v_\circ^\circ)](v_\circ^-) dv_+^- dv_\circ^-$$

where $\vartheta^t = \vartheta \cap \mathbb{X}^t$, $\mathcal{X}^t = \{\vartheta \in \mathcal{X} : \vartheta \subset \mathbb{X}^t\}$ and the sum is taken over all decompositions $\vartheta = v_\circ^- \sqcup v_\circ^\circ \sqcup v_+^\circ$ such that $v_\circ^\circ \in \mathcal{X}^t$ and $v_+^\circ \in \mathcal{X}^t$. The *multi-integrand* $\mathbf{M}(\mathbf{v})$ defines the values

$$\mathbf{M}(v_\cdot) = \delta_\emptyset(v_\cdot^+) \mathbf{M}(\mathbf{v}) \delta_\emptyset(v_\cdot^-), \quad \mathbf{v} = \begin{pmatrix} v_+^- & v_\circ^- \\ v_+^\circ & v_\circ^\circ \end{pmatrix}$$

of matrix elements $M(v) = M(v\cdot)$, $v = \sqcup_{\mu,\nu} v_\nu^\mu$, for a decomposable triangular tensor-operator $\mathbf{M} \equiv M$ defined on matrices $v\cdot = [v_\nu^\mu]_{\nu=-,\circ,+}^{\mu=-,\circ,+}$ whose elements are finite disjoint chains v_ν^μ , with $v_\nu^\mu = \emptyset$ for $\mu > \nu$ and with $M(v\cdot) = 0$ if $v\cdot^+ \neq \emptyset$ or $v\cdot^- \neq \emptyset$. The other values of $M(v\cdot)$ are defined by a kernel-operator function $M(\mathbf{v})$ of the quadruple $\mathbf{v} = (v_\nu^\mu)_{\nu=+,\circ}^{\mu=-,\circ}$ with continuous operator values $M(\mathbf{v}) : G_+ \rightarrow G_-$ in the scalar case $\mathfrak{k}_x = \mathbb{C}$. In the general case it is defined almost everywhere by its values on the chains $v_\nu^\mu \in \mathcal{X}$ in the continuous operators

$$M \begin{pmatrix} v_+^- & v_\circ^- \\ v_+^\circ & v_\circ^\circ \end{pmatrix} : \mathfrak{k}_{v_\circ^-}^\otimes \otimes \mathfrak{k}_{v_\circ^\circ}^\otimes \otimes G_+ \rightarrow \mathfrak{k}_{v_\circ^\circ}^\otimes \otimes \mathfrak{k}_{v_+^\circ}^\otimes \otimes G_-.$$

We will assume that these operators are bounded from $G_\star(q)$ to $G_\star(q^{-1})$ for some $q \in \mathfrak{p}_1$, such that $\|I\|_q = 1$, and that there exist strictly positive functions $r > 0$, $r^{-1} \in \mathfrak{p}_0$, and $s > 0$, $s^{-1} \in \mathfrak{p}_0$ such that

$$(5.2) \quad \|M\|_{q,t}^s(r) = \int_{\mathcal{X}^t} \|M_+^-(v)\|_{q,t}^s(r) dv < \infty, \quad \forall t < \infty,$$

where

$$\|M_+^-(v_+^-)\|_{q,t}^s(r) = \left(\int_{\mathcal{X}^t} \int_{\mathcal{X}^t} \operatorname{ess\,sup}_{v_\circ^\circ \in \mathcal{X}^t} (s(v_\circ^\circ) \|M(\mathbf{v})\|_q)^2 r(v_+^\circ \sqcup v_\circ^-) dv_+^\circ dv_\circ^- \right)^{\frac{1}{2}},$$

and $s(v) = \prod_{x \in v} s(x)$, $r(v) = \prod_{x \in v} r(x)$.

We mention that the single integral (3.2) corresponds to the case

$$M(v\cdot) = 0, \quad \forall v\cdot : \sum_{\mu \neq +, \nu \neq -} |v_\nu^\mu| \neq 1,$$

and $M(\mathbf{x}_\nu^\mu) = D_\nu^\mu(x)$ otherwise, where $\mathbf{x}_\nu^\mu$ denotes one of six ‘atomic’ triangular matrices $v\cdot(\mathbf{x}) = [v_\lambda^\kappa(\mathbf{x})]_{\lambda=-,\circ,+}^{\kappa=-,\circ,+} \equiv \mathbf{x}$ having all matrix elements $v_\lambda^\kappa(\mathbf{x})$ empty if $\mathbf{x} \neq \mathbf{x}_\lambda^\kappa$, but $v_\nu^\mu(\mathbf{x}) = x$ for $\mathbf{x} = \mathbf{x}_\nu^\mu$. Note that integrand $M(\mathbf{x}_\nu^\mu)$ is zero on the atomic matrices $\mathbf{x}_\nu^+$ and $\mathbf{x}_-^\mu$, otherwise $M(\mathbf{x}) = M(x)$, given by the single-point kernel $M(\mathbf{x})$ as a function of one of the four single-point tables $\mathbf{v}(x) = (v_\lambda^\kappa(\mathbf{x}))_{\lambda=+,\circ}^{\kappa=-,\circ} \equiv \mathbf{x}$:

$$(5.3) \quad \mathbf{x}_+^- = \begin{pmatrix} x & \emptyset \\ \emptyset & \emptyset \end{pmatrix}, \mathbf{x}_+^\circ = \begin{pmatrix} \emptyset & \emptyset \\ x & \emptyset \end{pmatrix}, \mathbf{x}_\circ^- = \begin{pmatrix} \emptyset & x \\ \emptyset & \emptyset \end{pmatrix}, \mathbf{x}_\circ^\circ = \begin{pmatrix} \emptyset & \emptyset \\ \emptyset & x \end{pmatrix},$$

determined by an $x \in \mathbb{X}$. It follows from the next theorem that the function $M(\mathbf{v})$ in (5.1) can be defined up to equivalence, whose kernel $\{M \approx 0\}$ consists of all multiple integrands with $\|M\|_{q,t}^s(r) = 0$ for all $t \in \mathbb{R}_+$ and for some q, r, s . In particular, M can be defined almost everywhere only for the tables $\mathbf{v} = (v_\nu^\mu)$ that give disjoint decompositions $v = \sqcup_{\mu,\nu} v_\nu^\mu$ of the chains $v \in \mathcal{X}$, that is, it may have nonzero values only on $\mathbf{v}$ representable in the form $\mathbf{v} = \sqcup_{x \in v} \mathbf{x}$, where $\mathbf{x}$ is one of the atomic tables given by (5.3) with indices μ, ν for $x \in v_\nu^\mu$.

Theorem 1. *Suppose that $M(\mathbf{v})$ is a locally integrable function in the sense of (5.2) for some $q, r, s > 0$. Then its integral (5.1) is a continuous operator $T_t = \mathfrak{I}_0^t(M)$*

from G_+ to G_- satisfying the estimate

$$(5.4) \quad \|T_t\|_p = \sup_{\chi \in G_*(p)} \{ \|T_t \chi\| (p^{-1}) / \|\chi\|(p) \} \leq \|M\|_{q,t}^s(r)$$

for any $p \geq r^{-1} + q + s^{-1}$. The operator T_t^* , formally adjoint to T_t in G_* , is the integral

$$(5.5) \quad \mathbf{i}_0^t(M)^* = \mathbf{i}_0^t(M^\dagger), \quad M^\dagger(v \cdot) = M(\widetilde{v \cdot})^*, \quad \widetilde{[v_\nu^\mu]} = [v_{-\nu}^{-\mu}]$$

of $\delta_0(v_+^+) M^*(\mathbf{v}) \delta_0(v_-) = M^\dagger(v \cdot)$, which is continuous from G_+ to G_- , and satisfying $\|M^*\|_{q,t}^s(r) = \|M\|_{q,t}^s(r)$ for $M^*(\mathbf{v}) = M(\mathbf{v}')^*$, where $(v_\nu^\mu)' = (v_{-\mu}^{-\nu})$. Moreover, the operator-valued function $t \mapsto T_t$ has the quantum-stochastic differential $dT_t = d\mathbf{i}_0^t(\mathbf{D})$ in the sense that

$$(5.6) \quad \mathbf{i}_0^t(M) = M(\emptyset) + \mathbf{i}_0^t(\mathbf{D}), \quad D_\nu^\mu(x) = \mathbf{i}_0^{t(x)}(\dot{M}(\mathbf{x}_\nu^\mu)),$$

defined by the quantum-stochastic derivatives $\mathbf{D} = [D_\nu^\mu]$ with values (3.1) acting from $G_*(p)$ to $G_*(p^{-1})$ and bounded almost everywhere:

$$\|D_+^-\|_{p,t}^{(1)} \leq \|M\|_{q,t}^s(r), \quad \|D\|_{p,t}^{(2)}(r) \leq \|M\|_{q,t}^s(r), \quad \|D_\circ^\circ\|_{p,t}^{(\infty)}(s) \leq \|M\|_{q,t}^s(r)$$

for $D = D_\circ^-$ and $D = D_+^\circ$, $p \geq r^{-1} + q + s^{-1}$. This differential is defined in the form of the multiple integrals (5.1), with respect to $\mathbf{v}$, of the point derivatives $\dot{M}(\mathbf{x}, v \cdot) = M(v \cdot \sqcup \mathbf{x})$, where $\mathbf{x}$ is given by one of four atomic tables (5.3) at a fixed point $x \in \mathbb{X}$.

Proof. Using property (4.3) in the form

$$\int \sum_{\sqcup v_\nu^\circ = \vartheta} f(v_-^\circ, v_\circ^\circ, v_+^\circ) d\vartheta = \iiint f(v_-^\circ, v_\circ^\circ, v_+^\circ) \prod_\nu dv_\nu^\circ,$$

it is easy to find that from the definition (5.1) for $\chi \in G_+$ we have

$$\begin{aligned} & \int \langle \chi(\vartheta) | [T_t \chi](\vartheta) \rangle d\vartheta \\ &= \int_{\mathcal{X}^t} dv_+^- \int_{\mathcal{X}^t} dv_+^\circ \int_{\mathcal{X}^t} dv_\circ^- \int_{\mathcal{X}^t} dv_\circ^\circ \langle \check{\chi}(v_\circ^\circ \sqcup v_+^\circ) | M(\mathbf{v}) \check{\chi}(v_\circ^- \sqcup v_\circ^\circ) \rangle \\ &= \int_{\mathcal{X}^t} dv_+^- \int_{\mathcal{X}^t} dv_+^\circ \int_{\mathcal{X}^t} dv_\circ^- \int_{\mathcal{X}^t} dv_\circ^\circ \langle M(\mathbf{v})^* \check{\chi}(v_\circ^\circ \sqcup v_+^\circ) | \check{\chi}(v_\circ^- \sqcup v_\circ^\circ) \rangle \\ &= \int \langle [T_t^* \chi](\vartheta) | \chi(\vartheta) \rangle d\vartheta, \end{aligned}$$

that is, T_t^* acts as $\mathbf{i}_0^t(M^*)$ in (5.1) with $M^*(\mathbf{v}) = M(\mathbf{v}')^*$, where $(v_\nu^\mu)' = (v_{-\mu}^{-\nu})$ with respect to the inversion $- : (-, \circ, +) \mapsto (+, \circ, -)$. More precisely, this yields $\|\mathbf{i}_0^t(M)\|_p = \|\mathbf{i}_0^t(M^*)\|_p$, since $\|T\|_p = \|T^*\|_p$ by the definition (5.4) of p -norm and by

$$\sup \{ |\langle \chi | T\chi \rangle| / \|\chi\|(p) \|\chi\|(p) \} = \sup \{ |\langle T^* \chi | \chi \rangle| / \|\chi\|(p) \|\chi\|(p) \}.$$

We estimate the integral $\langle \chi | T_t \chi \rangle$ using the Schwartz inequality

$$\int \|\check{\chi}(v)\|(q) \|\check{\chi}(v)\|(q) s^{-1}(v) dv \leq \|\check{\chi}\|(s^{-1}, q) \|\check{\chi}\|(s^{-1}, q)$$

and the property (4.3) of the multiple integral according to which

$$\|\dot{\chi}\|(s^{-1}, q) = \|\chi\|(q + s^{-1})$$

then

$$\begin{aligned} |\langle \chi | T_t \chi \rangle| &\leq \int_{\mathcal{X}^t} \int_{\mathcal{X}^t} \int_{\mathcal{X}^t} \int_{\mathcal{X}^t} \|\dot{\chi}(v_\circ^\circ \sqcup v_+^\circ)\|(q) \|M(\mathbf{v}) \dot{\chi}(v_\circ^\circ \sqcup v_\circ^-)\|(q^{-1}) d^4 \mathbf{v} \\ &\leq \int_{\mathcal{X}^t} dv_\circ^\circ \int_{\mathcal{X}^t} \int_{\mathcal{X}^t} \|\dot{\chi}(v_\circ^\circ \sqcup v_+^\circ)\|(q) \left(\int_{\mathcal{X}^t} \|M(\mathbf{v})\|_q dv_+^- \right) \|\dot{\chi}(v_\circ^- \sqcup v_\circ^\circ)\|(q) dv_\circ^- dv_+^\circ \\ &\leq \int_{\mathcal{X}^t} dv \left(\int_{\mathcal{X}^t} \|\dot{\chi}(v \sqcup v_+^\circ)\|^2(q) \frac{dv_+^\circ}{r(v_+^\circ)} \int_{\mathcal{X}^t} \|\dot{\chi}(v \sqcup v_\circ^-)\|^2(q) \frac{dv_\circ^-}{r(v_\circ^-)} \right)^{\frac{1}{2}} \|M_\circ^\circ(v)\|_{q,t}(r) \\ &= \int_{\mathcal{X}^t} dv \|\dot{\chi}(v)\|(r^{-1} + q) \|M_\circ^\circ(v)\|_{q,t}(r) \|\dot{\chi}(v)\|(r^{-1} + q) \\ &\leq \operatorname{ess\,sup}_{v \in \mathcal{X}^t} \left\{ s(v) \|M_\circ^\circ(v)\|_{q,t}(r) \right\} \|\chi\|(r^{-1} + q + s^{-1}) \|\chi\|(r^{-1} + q + s^{-1}), \end{aligned}$$

where $\|M_\circ^\circ(v_\circ^\circ)\|_{q,t}(r) = (\int_{\mathcal{X}^t} \int_{\mathcal{X}^t} (\int_{\mathcal{X}^t} \|M(\mathbf{v})\|_q dv_+^-)^2 r(v_\circ^- \sqcup v_+^\circ) dv_\circ^- dv_+^\circ)^{1/2}$. Then since

$$\operatorname{ess\,sup}_{v \in \mathcal{X}^t} \{s(v) \|M_\circ^\circ(v)\|_{q,t}(r)\} \leq \|M\|_{q,t}^s(r),$$

and since

$$\sup_{\chi \in G_\star(p)} \frac{\|T\chi\|(p^{-1})}{\|\chi\|(p)} = \sup_{\chi \in G_\star(p)} \frac{|\langle \chi | T\chi \rangle|}{\|\chi\|(p) \|\chi\|(p)},$$

it follows that $\|T_t\|_p \leq \|M\|_{q,t}^s(r)$ for all $p \geq r^{-1} + q + s^{-1}$.

Now using the definition (5.1) and the property

$$\int_{\mathcal{X}^t} \chi(\vartheta) d\vartheta = \chi(\emptyset) + \int_{\mathbb{X}^t} dx \int_{\mathcal{X}^{t(x)}} \dot{\chi}(x, \vartheta) d\vartheta,$$

where $\dot{\chi}(x, \vartheta) = \chi(\vartheta \sqcup x)$, it is easy to see that $[(T_t - T_0)\chi](\vartheta) = [(\mathbf{t}_0^t(M) - M(\emptyset))\chi](\vartheta) =$

$$\begin{aligned} &= \int_{\mathbb{X}^t} dx \sum_{\substack{t(v_\nu^\circ) < t(x) \\ v_\circ^\circ \sqcup v_+^\circ \subseteq \vartheta}} \left\{ \int_{\mathcal{X}^{t(x)}} dv_+^- \left[\int_{\mathcal{X}^{t(s)}} dv_\circ^- (\dot{M}(\mathbf{x}_+^-, \mathbf{v}) \dot{\chi}(v_\circ^- \sqcup v_\circ^\circ) \right. \right. \\ &\quad \left. \left. + \dot{M}(\mathbf{x}_\circ^-, \mathbf{v}) \dot{\chi}(x \sqcup v_\circ^- \sqcup v_\circ^\circ) \right) \right] (\vartheta \setminus v_\circ^\circ \setminus v_+^\circ) \right. \\ &\quad \left. + \sum_{x \in \mathbb{X}^t} \int_{\substack{t(v_\nu^\circ) < t(x) \\ v_\circ^\circ \sqcup v_+^\circ \subseteq \vartheta}} \left\{ \int_{\mathcal{X}^{t(x)}} dv_+^- \left[\int_{\mathcal{X}^{t(x)}} dv_\circ^- (\dot{M}(\mathbf{x}_+^\circ, \mathbf{v}) \dot{\chi}(v_\circ^- \sqcup v_\circ^\circ) \right. \right. \right. \\ &\quad \left. \left. + \dot{M}(\mathbf{x}_\circ^\circ, \mathbf{v}) \dot{\chi}(x \sqcup v_\circ^- \sqcup v_\circ^\circ) \right) \right] ((\vartheta \setminus v_\circ^\circ \setminus v_+^\circ) \right. \\ &= \int_{\mathbb{X}^t} dx [D_+^-(x)\chi + D_\circ^-(x)\dot{\chi}(x)](\vartheta) + \sum_{x \in \mathbb{X}^t} [D_+^\circ(x)\chi + D_\circ^\circ(x)\dot{\chi}(x)](\vartheta \setminus x). \end{aligned}$$

Consequently, $T_t - T_0 = \sum \Lambda_\nu^\mu(D_\nu^\mu, \mathbb{X}^t)$, where $\Lambda_\nu^\mu(D, \Delta)$ are defined in (3.3) as operator-valued measures on $\mathbb{X}$ of operator-functions

$$\begin{aligned} [D_+^\mu(x)\chi](\vartheta) &= \sum_{v_\circ^\circ \sqcup v_+^\circ \subseteq \vartheta}^{t(v_\nu^\circ) < t(x)} \int_{\mathcal{X}^{t(x)}} dv_+^- \int_{\mathcal{X}^{t(x)}} dv_\circ^- [\dot{M}(\mathbf{x}_+^\mu, \mathbf{v}) \dot{\chi}(v_\circ^- \sqcup v_+^\circ)](\vartheta_-^\circ), \\ [D_\circ^\mu(x)\dot{\chi}(x)](\vartheta) &= \sum_{v_\circ^\circ \sqcup v_+^\circ \subseteq \vartheta}^{t(v_\nu^\circ) < t(x)} \int_{\mathcal{X}^{t(x)}} dv_+^- \int_{\mathcal{X}^{t(x)}} dv_\circ^- [\dot{M}(\mathbf{x}_\circ^\mu, \mathbf{v}) \dot{\chi}(x \sqcup v_\circ^- \sqcup v_+^\circ)](\vartheta_-^\circ), \end{aligned}$$

acting on $\chi \in G_+$ and $\dot{\chi}(v) \in \mathfrak{k}_v^\otimes \otimes G_+$, where $\vartheta_-^\circ = \vartheta \cap \overline{(v_\circ^\circ \sqcup v_+^\circ)} = \vartheta \setminus v_\circ^\circ \setminus v_+^\circ$. This can be written in terms of (5.1) as

$$D_\nu^\mu(x) = \mathfrak{z}_0^t(\dot{M}(\mathbf{x}_\nu^\mu)).$$

Because of the inequality $\|T_t\|_p \leq \|M\|_{q,t}^s(r)$ for all $p \geq r^{-1} + q + s^{-1}$ we obtain $\|D_+^\mu\|_{p,t}^{(1)} \leq \|M\|_{q,t}^s(r)$, since $\|D_+^\mu(x)\|_p \leq \|\dot{M}(\mathbf{x}_+^\mu)\|_{q,t(x)}^s(r)$:

$$\begin{aligned} \int_{\mathbb{X}^t} \|D_+^\mu(x)\|_p dx &\leq \int_{\mathbb{X}^t} \|\dot{M}(\mathbf{x}_+^\mu)\|_{q,t(x)}^s(r) dx = \int_{\mathbb{X}^t} dx \int_{\mathcal{X}^{t(x)}} \|M_+^\mu(x \sqcup v)\|_{q,t(x)}^s(r) dv \\ &= \int_{\mathcal{X}^t} \|M_+^\mu(v)\|_{q,t}^s(r) dv - \|M_+^\mu(\emptyset)\|_{q,t}^s(r) \\ &= \|M\|_{q,t}^s(r) - \|M_+^\mu(\emptyset)\|_{q,t}^s(r). \end{aligned}$$

For the estimate of D_+° we shall require the use of the norm

$$\|M_+^\circ\|_{q,t}^s(r) := \left(\int_{\mathcal{X}^t} \left(\|M_+^\circ(v_+^\circ)\|_{q,t}^s(r) \right)^2 r^\otimes(v_+^\circ) dv_+^\circ \right)^{1/2},$$

where

$$\|M_+^\circ(v_+^\circ)\|_{q,t}^s(r) := \left(\int_{\mathcal{X}^t} \left(\int_{\mathcal{X}^t} \operatorname{ess\,sup}_{v_\circ^\circ \in \mathcal{X}^t} s(v_\circ^\circ) \|M(\mathbf{v})\|_q dv_+^- \right)^2 r^\otimes(v_\circ^\circ) dv_\circ^- \right)^{1/2};$$

in particular, $\|M_+^\circ\|_{q,t}^s(r) \leq \|M\|_{q,t}^s(r)$. So we have

$$\begin{aligned} \|D_+^\circ\|_{p,t}^{(2)}(r)^2 &\leq \int_{\mathbb{X}^t} (\|\dot{M}_+^\circ(\mathbf{x}_+^\circ)\|_{q,t(x)}^s(r))^2 r(x) dx \\ &= \|M_+^\circ\|_{q,t}^s(r)^2 - \|M_+^\circ(\emptyset)\|_{q,t}^s(r)^2 \leq \|M\|_{q,t}^s(r)^2. \end{aligned}$$

In a similar manner we obtain the estimate for $D_\circ^-$,

$$\begin{aligned} \|D_\circ^-\|_{p,t}^{(2)}(r)^2 &\leq \int_{\mathbb{X}^t} (\|\dot{M}_\circ^-(\mathbf{x}_\circ^-)\|_{q,t(x)}^s(r))^2 r(x) dx \\ &= \|M_\circ^-\|_{q,t}^s(r)^2 - \|M_\circ^-(\emptyset)\|_{q,t}^s(r)^2 \leq \|M\|_{q,t}^s(r)^2. \end{aligned}$$

Finally, from $\|D_\circ^\circ(x)\|_p \leq \|\dot{M}(\mathbf{x}_\circ^\circ)\|_{q,t(x)}^s(r)$ we similarly obtain

$$\|D_\circ^\circ\|_{p,t}^{(\infty)}(s) \leq \operatorname{ess\,sup}_{x \in \mathbb{X}^t} \{s(x) \|\dot{M}(\mathbf{x}_\circ^\circ)\|_{q,t(x)}^s(r)\} \leq \|M\|_{q,t}^s(r)$$

if $p \geq r^{-1} + q + s^{-1}$, which concludes the proof. $\square$

Since $T_t = \mathfrak{I}_0^t(M)$ the theorem obviously states that $\|\mathfrak{I}_0^t(M)\|_p \leq \|M\|_{q,t}^s(r)$ for all $p \geq \frac{1}{r} + q + \frac{1}{s}$, and in particular for the case when $M(\mathfrak{v}) = 0$ if $\sum |v_\nu^\mu| \neq 1$, and $M(\mathbf{x}) = D(\mathbf{x})$. Then the result of the theorem becomes

$$\|\mathfrak{I}_0^t(D)\|_p \leq \|D\|_{q,t}^s(r)$$

and now we shall begin to evaluate the quantum stochastic norm (5.2) explicitly for the single integrand $D(\mathbf{x})$.

$$\|D\|_{q,t}^s(r) = \|D_+^-\|_{q,t}^{(1)} + \|D_+^-(\emptyset)\|_{q,t}^s(r)$$

one may now proceed in a similar manner to find that

$$\|D_+^-(\emptyset)\|_{q,t}^s(r) = \|D_\circ^-\|_{q,t}^{(2)}(r) + \|D_+^\circ\|_{q,t}^{(2)}(r) + \|D_\circ^\circ\|_{q,t}^{(\infty)}(s),$$

and thus we have recovered that inequality stated at the beginning of this section.

6. ADAPTED AND Q-ADAPTED QS INTEGRALS

The quantum-stochastic integral (5.1) constructed in [7], as well as its single variations (3.2) introduced in [6], are defined explicitly and do not require that the functions M and D under the integral be adapted. By virtue of the continuity we have proved above, they can be approximated in the inductive convergence by the sequence of integral sums $\mathfrak{I}_0^t(M_n)$, $\mathfrak{I}_0^t(D_n)$ corresponding to step measurable operator functions M_n and D_n if the latter converge inductively to M and D in the polynomial norm (5.2). In fact, if there exist functions r, s with $r^{-1}, s^{-1} \in \mathfrak{p}_0$ and $q \in \mathfrak{p}_1$ such that $\|M_n - M\|_{q,t}^s(r) \rightarrow 0$, then there also exists a function $p \in \mathfrak{p}_1$ such that $\|\mathfrak{I}_0^t(M_n - M)\|_p \rightarrow 0$, and we have $p \geq r^{-1} + q + s^{-1}$ by the inequality (5.4), which implies the inductive convergence $\mathfrak{I}_0^t(M_n) \rightarrow \mathfrak{I}_0^t(M)$ as a result of the linearity of $\mathfrak{I}_0^t$.

Let $Q : x \mapsto \mathfrak{L}(\mathfrak{k}_x)$ be a measurable adjointable operator-valued function with q -contractive values $\|Q(x)\| \leq q(x)$ with respect to a positive function $q \in \mathfrak{p}$. We shall say that the integrand $D(x)$ is Q -adapted if it has the product form

$$(6.1) \quad D_\nu^\mu(x) = D_\nu^\mu(x)^{t(x)} \otimes Q_{t(x)}^\otimes$$

with respect to the Fock split $\mathcal{G}_* = \mathcal{G}_*^{t(x)} \otimes \mathcal{F}_{*[t(x)]}$ corresponding to $\mathbb{X} = \mathbb{X}^{t(x)} \sqcup \mathbb{X}_{[t(x)]}$, where $D_\nu^\mu(x)^{t(x)} \equiv K_\nu^\mu(x)$ is the restricted action $G_+^{t(x)} \rightarrow G_-^{t(x)}$ of $D_\nu^\mu(x)$ corresponding to the projections $P_{t(x)} D_\nu^\mu(x) P_{t(x)}$ where the projection P_t is given on $\mathcal{G}_*$ by

$$[P_t \chi](\vartheta) = \chi(\vartheta^t) \delta_\emptyset(\mathfrak{x}), \quad P_0 \equiv P_\emptyset$$

where $\mathfrak{x} = \vartheta \setminus \vartheta^t$, and $\delta_\emptyset(\vartheta) = 1$ if ϑ is empty, and is otherwise zero. We can now write the Q -adapted QS integrals as a more explicit form of the general QS integrals

(3.3)

$$\begin{aligned}
[\Lambda_-^+(\mathbf{D}_+^-, \Delta)\chi](\vartheta) &= \int_{\Delta} [K_+^-(x) \otimes Q^{\otimes}(\kappa)\dot{\chi}(\kappa)](\vartheta^{t(x)})dx, \quad x \notin \vartheta \\
[\Lambda_+^+(\mathbf{D}_+^{\circ}, \Delta)\chi](\vartheta) &= \sum_{x \in \Delta \cap \vartheta} [K_+^{\circ}(x) \otimes Q^{\otimes}(\kappa)\dot{\chi}(\kappa)](\vartheta^{t(x)}), \\
[\Lambda_-^{\circ}(\mathbf{D}_+^-, \Delta)\chi](\vartheta) &= \int_{\Delta} [K_+^-(x) \otimes Q^{\otimes}(\kappa)\dot{\chi}(x \sqcup \kappa)](\vartheta^{t(x)})dx, \quad x \notin \vartheta \\
(6.2) \quad [\Lambda_+^{\circ}(\mathbf{D}_+^{\circ}, \Delta)\chi](\vartheta) &= \sum_{x \in \Delta \cap \vartheta} [K_+^{\circ}(x) \otimes Q^{\otimes}(\kappa)\dot{\chi}(x \sqcup \kappa)](\vartheta^{t(x)}).
\end{aligned}$$

Here $Q^{\otimes}(\kappa) = \otimes_{x \in \kappa} Q(x)$ for any $\kappa = \vartheta_{t(x)} \in \mathcal{X}_{t(x)}$ and $\dot{\chi}(\kappa, \vartheta^t) = \chi(\vartheta^t \sqcup \kappa)$.

In the next proposition we look at the some of the properties of a special class of Q -adapted integrals. In particular, we consider the g -commutator of the creation and annihilation integrals $\Lambda(\mathbf{D}_+^{\circ})$ and $\Lambda(\mathbf{D}_+^{-})$, given by

$$(6.3) \quad [\Lambda(\mathbf{D}_+^{-}), \Lambda(\mathbf{D}_+^{\circ})]_g := \Lambda(\mathbf{D}_+^{-})\Lambda(\mathbf{D}_+^{\circ}) - g(\Lambda(\mathbf{D}_+^{\circ})\Lambda(\mathbf{D}_+^{-})),$$

where g is considered to be an element of $L^{\infty}(\mathbb{X} \times \mathbb{X})$ in the sense that its action on $\Lambda(\mathbf{D}_+^{\circ})\Lambda(\mathbf{D}_+^{-})$ is defined as

$$(6.4) \quad g(\Lambda(\mathbf{D}_+^{\circ})\Lambda(\mathbf{D}_+^{-})) = \iint \Lambda_+^+(dz)D_+^{\circ}(z)g(z, x)D_+^{-}(x)\Lambda_+^{\circ}(dx),$$

with respect to the identifications

$$\Lambda(\mathbf{D}_+^{\circ}(z), dz) = \Lambda_+^+(dz)D_+^{\circ}(z), \quad \Lambda(\mathbf{D}_+^{-}(x), dx) = D_+^{-}(x)\Lambda_+^{\circ}(dx),$$

$x, z \in \mathbb{X}$.

Proposition 1. *Let $\Lambda(\mathbf{D}_+^{-})$ and $\Lambda(\mathbf{D}_+^{\circ})$ be Q -adapted quantum stochastic integrals of the form*

$$\begin{aligned}
[\Lambda(\mathbf{D}_+^{-}, \mathbb{X}^t)\chi](\vartheta) &= \int_{\mathbb{X}^t} K_+^-(x, \vartheta^{t(x)}) \otimes Q^{*\otimes}(\vartheta_{t(x)})\chi(\vartheta \sqcup x)dx, \\
[\Lambda(\mathbf{D}_+^{\circ}, \mathbb{X}^t)\chi](\vartheta) &= \sum_{z \in \vartheta^t} K_+^{\circ}(z, \vartheta^{t(z)}) \otimes Q^{\otimes}(\vartheta_{t(z)})\chi(\vartheta \setminus z),
\end{aligned}$$

such that $K_+^-(x)$ and $K_+^{\circ}(x)$ are diagonal on $\mathcal{X}^{t(x)}$ in the sense that $[K_+^{\circ}(x)\chi](\vartheta^{t(x)}) = K_+^{\circ}(x, \vartheta^{t(x)})\chi(\vartheta^{t(x)})$ and $[K_+^-(x)\dot{\chi}](\vartheta^{t(x)}) = K_+^-(x, \vartheta^{t(x)})\chi(x \sqcup \vartheta^{t(x)})$, where Q is q -contractive normal operator $[Q, Q^*] = 0$. Further, suppose that the operator $K_+^-(x)$ commutes with $K_+^{\circ}(z)$ in $\mathcal{G}_*$ such that $[K_+^-(x), K_+^{\circ}(z)] = 0$ for all $x \neq z \in \mathbb{X}^t$, and that $[K_+^-(x), Q^{\otimes}] = 0 = [K_+^{\circ}(x), Q^{\otimes}]$ in $\mathcal{G}_*$, then the g -commutator (6.3) satisfies the equation

$$(6.5) \quad [\Lambda(\mathbf{D}_+^{-}, \mathbb{X}^t), \Lambda(\mathbf{D}_+^{\circ}, \mathbb{X}^t)]_g = \int_{\mathbb{X}^t} K_+^-(x)K_+^{\circ}(x)dx \otimes (Q^*Q)^{\otimes}$$

in $\mathcal{G}_*^t \otimes \mathcal{F}_{*t}$ if we have $K_+^-(x, \vartheta^{t(x)})Q(x) = g(z, x)K_+^-(x, \vartheta^{t(x)} \setminus z) \otimes I_z$ for all $z \in \vartheta^{t(x)}$, and $Q^*(z)K_+^{\circ}(z, \vartheta^{t(z)} \sqcup x) = I_x \otimes K_+^{\circ}(z, \vartheta^{t(z)})g(z, x)$ for all $x < z$ with $x \notin \vartheta^{t(z)}$.

Proof. First we calculate

$$\begin{aligned} & [\Lambda(\mathbf{D}_\circ^-, \mathbb{X}^t) \Lambda(\mathbf{D}_+^\circ, \mathbb{X}^t) \chi](\vartheta) - \left[\int_{\mathbb{X}^t} K_\circ^-(x) K_+^\circ(x) dx \otimes (Q^* Q)^\otimes \chi \right](\vartheta) = \\ &= \int_{\mathbb{X}^t} \sum_{z \in \vartheta^{t(x)}} K_\circ^-(x, \vartheta^{t(x)}) K_+^\circ(z, \vartheta^{t(z)}) Q^\otimes(\vartheta_{t(z)}^{t(x)} \sqcup x) (Q^* Q)^\otimes(\vartheta_{t(x)}) \chi(x \sqcup \vartheta \setminus z) dx \\ &+ \sum_{z \in \vartheta} \int_{\mathbb{X}^{t(z)}} K_\circ^-(x, \vartheta^{t(x)}) Q^{*\otimes}(\vartheta_{t(x)}^{t(z)} \sqcup z) K_+^\circ(z, \vartheta^{t(z)} \sqcup x) (Q^* Q)^\otimes(\vartheta_{t(z)}) \chi(x \sqcup \vartheta \setminus z) dx, \end{aligned}$$

where $\vartheta_s^t = \vartheta \cap \mathbb{X}_s^t$, and then we calculate $g[\Lambda(\mathbf{D}_+^\circ, \mathbb{X}^t) \Lambda(\mathbf{D}_\circ^-, \mathbb{X}^t) \chi](\vartheta) =$

$$\begin{aligned} &= \int_{\mathbb{X}^t} \sum_{z \in \vartheta^{t(x)}} g(z, x) K_+^\circ(z, \vartheta^{t(z)}) Q^\otimes(\vartheta_{t(z)}^{t(x)}) K_\circ^-(x, \vartheta^{t(x)} \setminus z) (Q^* Q)^\otimes(\vartheta_{t(x)}) \chi(x \sqcup \vartheta \setminus z) dx \\ &+ \sum_{z \in \vartheta} \int_{\mathbb{X}^{t(z)}} g(z, x) K_+^\circ(z, \vartheta^{t(z)}) K_\circ^-(x, \vartheta^{t(x)}) Q^{*\otimes}(\vartheta_{t(x)}^{t(z)}) (Q^* Q)^\otimes(\vartheta_{t(z)}) \chi(x \sqcup \vartheta \setminus z) dx, \end{aligned}$$

thus when equating the integrands for these two formula, using the commutativity of the factors, we find that for $z \in \vartheta^{t(x)}$

$$K_\circ^-(x, \vartheta^{t(x)}) Q(x) = g(z, x) K_\circ^-(x, \vartheta^{t(x)} \setminus z)$$

and for $x < z$ we find

$$Q^*(z) K_+^\circ(z, \vartheta^{t(z)} \sqcup x) = K_+^\circ(z, \vartheta^{t(z)}) g(z, x).$$

□

Notice that when $g(z, x)$ is separable we have $g(z, x) I_x = h(z) Q(x)$ for $z < x$, and $g(z, x) I_z = Q^*(z) f(x)$ when $z > x$, and also note that if $h = f^*$ then g is self-adjoint, $g^*(x, z) = g(z, x)$, having the form

$$(6.6) \quad g(z, x) = h^*(z) q(x) 1_{t(x)}(z) + q^*(z) h(x) 1_{t(z)}(x), \quad z \neq x$$

where $1_{t(x)}(z) = 1$ if $z < x$ otherwise 0. Further, from this separability of g it follows that $K_\circ^-(x, v) = k_\circ^-(x) \otimes h^\otimes(v) I_v$ for all $v \in \mathcal{X}^{t(x)}$, and that $K_+^\circ(z, v) = k_+^\circ(z) \otimes f^\otimes(v) I_v$ for all $v \in \mathcal{X}^{t(z)}$. Consider for example the case when $g(z, x) = \exp\{ip(x - z)\}$, where $p : \mathbb{X} \rightarrow \mathbb{R}$ is linear functional $p(x - z) = p(x) - p(z)$.

Corollary 1. *Suppose that $g(x, z)$, $x \neq z$, is a constant $\mathbb{R}$ -valued function of the form*

$$g(z, x) = 1(z) q(x) 1_{t(x)}(z) + q(z) 1(x) 1_{t(z)}(x), \quad q(x) = q \in \mathbb{R} \forall x$$

where $1_t(x) = 1$ if $t(x) < t$ and otherwise zero, then we define the q -commutator

$$(6.7) \quad [\Lambda(\mathbf{D}_\circ^-), \Lambda(\mathbf{D}_+^\circ)]_q = \Lambda(\mathbf{D}_\circ^-) \Lambda(\mathbf{D}_+^\circ) - q \Lambda(\mathbf{D}_+^\circ) \Lambda(\mathbf{D}_\circ^-),$$

and we recover the commutation relations of the Bosonic and Fermionic quantum fields under the above conditions with $Q(x) = qI$ when we respectively set $q = +1$ and $q = -1$, such that

$$[\Lambda(\mathbf{D}_\circ^-, \mathbb{X}^t), \Lambda(\mathbf{D}_+^\circ, \mathbb{X}^t)]_{\pm 1} = \int_{\mathbb{X}^t} K_\circ^-(x) K_+^\circ(x) dx \otimes I^\otimes$$

and thus we have the representations of the Bosonic and Fermionic quantum fields in the Guichardet-Fock space as $\pm I$ -adapted QS integrals.

Further, the trivial commutator of the monotonic quantum field is obtained when $Q = O$, corresponding also to $q = 0$, such that

$$[\Lambda(\mathbf{D}_\circ^-, \mathbb{X}^t), \Lambda(\mathbf{D}_+^\circ, \mathbb{X}^t)]_0 = \int_{\mathbb{X}^t} K_\circ^-(x) K_+^\circ(x) dx \otimes O^\otimes$$

where we have made use of the identification $D_\nu^\mu(\emptyset) = 0$. Such O -adapted monotonic calculus, also called the vacuum adapted calculus as $O^\otimes = P_\emptyset$, was also studied by Belton in [9].

Obviously the QS integral $X(t) = \mathbf{i}_0^t(\mathbf{D})$ of any Q -adapted integrand $\mathbf{D}(x)$ is an operator-valued Q -adapted process in the sense that $X(t) = X^t \otimes Q_t^\otimes$. The approximation of this integral in the class of adapted step functions, leads by continuity to the usual definition of the quantum-stochastic integral $\mathbf{i}_0^t(\mathbf{D})$ which was given by Hudson and Parthasarathy corresponding to the I -adapted case with $\mathbb{X} = \mathbb{R}_+$ such that $x \equiv t(x)$. That is the weak limit of integral sums

$$\mathbf{i}_0^t(\mathbf{D}_n) = \int_0^t \Lambda(\mathbf{D}_n, dx) = \sum_{j=1}^n D_\nu^\mu(x_j) A_\mu^\nu(\Delta_j),$$

where $\mathbf{D}(x_j) = \mathbf{D}_n(x)$ for $x \in [x_j, x_{j+1})$ is an adapted approximation corresponding to the decomposition $\mathbb{R}_+ = \sum_{j=1}^n \Delta_j$ into the intervals $\Delta_j = [x_j, x_{j+1})$ given by the chain $x_0 = 0 < x_1 < \dots < x_n < x_{n+1} = \infty$, and $D_\nu^\mu(x) A_\mu^\nu(\Delta)$ is the sum of the operators (3.3) with functions $D_\nu^\mu(x)$ constant on Δ which can therefore be pulled out in front of the integrals Λ_μ^ν such that $A_\mu^\nu(\Delta) = \int_\Delta \Lambda_\mu^\nu(dx)$.

In particular, for $D_+^- = 0 = D_\circ^\circ$ and $D_\circ^- = k \otimes \hat{1} = D_+^\circ$, where $\hat{1} = I^\otimes$ is the unit operator in $\mathcal{F}_*$ and $k(x)$ is a scalar locally square integrable function corresponding to the case $\mathfrak{k}_x = \mathbb{C} = \mathfrak{h}$, we obtain the Itô definition of the Wiener integral

$$\dot{I}_0^t(k) = \int_0^t k(x) w(dx), \quad \int_0^t k(x) \widehat{w}(dx) = \mathbf{i}_0^t(\mathbf{D})$$

with respect to the stochastic measure $w(\Delta)$, $\Delta \in \mathfrak{F}_\mathbb{X}$ on $\mathbb{R}_+$, represented in $\mathcal{G}_* = \mathcal{F}_*$ by the operators $\widehat{w}(\Delta) = A_\circ^+(\Delta) + A_\circ^-(\Delta)$. We also note that the multiple integral (5.1) in the trivially adapted case $M(\mathbf{v}) = M(\mathbf{v}) \otimes I^\otimes$ defines the Fock representation of the generalized Maassen-Meyer kernels [10], [17] and in the case

$$M(\mathbf{v}) = m(v_\circ^- \sqcup v_+^\circ) \delta_\emptyset(v_+^-) \delta_\emptyset(v_\circ^\circ), \quad \delta_\emptyset(v) = \begin{cases} 1, & v = \emptyset, \\ 0, & v \neq \emptyset \end{cases}$$

it leads to the multiple stochastic integrals $\mathbf{i}_0^t(M) = \widehat{I}_0^t(m)$,

$$I_0^t(m) = \sum_{n=0}^{\infty} \int_{0 \leq t_1 < \dots < t_n < t} \dots \int m(x_1, \dots, x_n) w(dx_1) \dots w(dx_n)$$

of the generalized functions $m \in \bigcup_{r^{-1} \in \mathfrak{p}_0} G_\star(r)$, that is, to the Hida distributions [12], [19] of the Wiener measure $w(\Delta)$ represented as $\widehat{w}(\Delta)$. Thus, we can consider the trivially adapted QS multiple integrals $\mathfrak{I}_0^t(M \otimes \mathbb{I}^\otimes)$ as quantum Hida operator-distributions whose properties are described in the following corollary when $\mathbb{Q} = \mathbb{I}$.

Corollary 2. *Suppose that $M(\mathbf{v}) = M(\mathbf{v}) \otimes \mathbb{Q}^\otimes$ where $\|\mathbb{Q}^\otimes\|_q = 1$, i.e. the operator-function M is defined by the q -contractive ampliation of the $\star$ -kernel M with $\|M\|_t^s(r) < \infty$, where*

$$M \begin{pmatrix} v_+^-, & v_\circ^- \\ v_+^\circ, & v_\circ^\circ \end{pmatrix} : \mathfrak{k}^\otimes(v_\circ^- \sqcup v_\circ^\circ) \otimes \mathfrak{h} \rightarrow \mathfrak{k}^\otimes(v_\circ^\circ \sqcup v_+^\circ) \otimes \mathfrak{h},$$

and $\|M\|_t^s(r) < \infty$ for all $t \in \mathbb{R}_+$, and for some r, s with $r^{-1}, s^{-1} \in \mathfrak{p}_0$, where

$$\|M\|_t^s(r) = \int_{\mathcal{X}^t} dv_+^- \left(\int_{\mathcal{X}^t} dv_+^\circ \int_{\mathcal{X}^t} dv_\circ^- \text{ess sup}_{v_\circ^\circ \in \mathcal{X}^t} \{s(v_\circ^\circ) \|M(\mathbf{v})\|\}^2 r(v_+^\circ \sqcup v_\circ^-) \right)^{1/2}$$

and $r(v) = \prod_{x \in v} r(x)$, $s(v) = \prod_{x \in v} s(x)$. Then the multiple integral (5.1) defines a $\mathbb{Q}$ -adapted family T_t , $t \in \mathbb{R}_+$, of p -bounded operators

$$T_t = \mathfrak{I}_0^t(M \otimes \mathbb{Q}^\otimes), \quad \|T_t\|_p \leq \|M\|_t^s(r)$$

for $p \geq r^{-1} + q + s^{-1}$, with bounded $\mathbb{Q}$ -adapted quantum-stochastic derivatives

$$D_\nu^\mu(x) = \mathfrak{I}_0^{t(x)}(\dot{M}(\mathbf{x}_\nu^\mu) \otimes \mathbb{Q}^\otimes) \equiv K_\nu^\mu(x) \otimes \mathbb{Q}_{t(x)}^\otimes.$$

Proof. Since $\|M(\mathbf{v}) \otimes \mathbb{Q}^\otimes\|_q \leq \|M(\mathbf{v})\|$ the result immediately follows from the inequality $\|\mathfrak{I}_0^t(M \otimes \mathbb{Q}^\otimes)\|_p \leq \|M \otimes \mathbb{Q}^\otimes\|_{q,t}^s(r) \leq \|M\|_t^s(r)$ for $p \geq r^{-1} + q + s^{-1}$. $\square$

REFERENCES

- [1] L. Accardi and F. Fagnola, *Stochastic Integration*, Vol. 1303, Lecture Notes in Mathematics, 1988.
- [2] L. Accardi and J. Quaegebeur, *The Itô Algebra of Quantum Gaussian Fields*, J. Funct. Anal., 1989, 85:213–263.
- [3] V. P. Belavkin, *Reconstruction Theorem for Quantum Stochastic Processes*, Theoretical Mathematical Physics, 1985, 3:409–431.
- [4] V. P. Belavkin, *Non-Demolition Measurements, Non-Linear Filtering, and Dynamic Programming of Quantum Stochastic Processes*, Proceedings of Bellman Continuum Workshop, 1988.
- [5] V. P. Belavkin, *Optimal Non-Linear Filtering of Quantum Signals*, Proceedings of 9th Conference on Coding Theory and Information Transmission, University of Odessa, 1988, 342–345.
- [6] V. P. Belavkin, *A Quantum Stochastic Calculus in Fock Space of Input and Output Non-Demolition Processes*, Vol. 1442, Lecture Notes in Mathematics, Springer-Verlag, 1990, 99–125.
- [7] V. P. Belavkin, *A Quantum Non-Adapted Itô Formula and Stochastic Analysis in Fock Scale*, J. Funct. Anal., 1991, 102:414–447.
- [8] V. P. Belavkin, *Quantum Stochastic Calculus and Quantum Non-Linear Filtering*, J. Multivariate Analysis, 1992, 42:171–201.
- [9] A. C. R. Belton, *A matrix formulation of quantum stochastic calculus*, PhD thesis, University of Oxford, 1998.

- [10] M. P. Evans and R. L. Hudson, *Multidimensional Quantum Diffusions*, Vol. 1303, Lecture Notes in Mathematics, 1988.
- [11] A. Guichardet, *Symmetric Hilbert Spaces and Related Topics*, Springer-Verlag, 1972.
- [12] T. Hida, *Brownian Motion*, Springer-Verlag, 1980.
- [13] A. S. Holevo, *Time-Ordered Exponentials in Quantum Stochastic Calculus*, Preprint 517, Universität Heidelberg, 1989.
- [14] J. M. Lindsay and H. Maassen, *The Stochastic Calculus of Bose Noise*, CWI syllabus 32, Centre for Mathematics and Computer science Amsterdam, 1992.
- [15] J. M. Lindsay, *On Set Convolutions and Integral-Sum Kernel Operators*, Proceedings Int. Conf. on Probability Theory and Mathematical Statistics, Vilnius, 1990.
- [16] P. Malliavin, *Stochastic Calculus of Variations and Hypocoelliptic Operators*, Proceedings of the International Symposium on Stochastic Differential Equations, New York, 1978, 195–293.
- [17] P. A. Meyer, *Éléments de Probabilités Quantiques*, Vol. 1247, Lecture Notes in Mathematics, 1987, 33–78.
- [18] K. R. Parthasarathy and K. B. Sinha, *Stochastic Integral Representation of Bounded Quantum Martingales in Fock Space*, J. Funct. Anal., 1986, 67:126–151.
- [19] Yu. G. Kondratiev, P. Leukert, J. Potthoff, L. Streit, and W. Westerkamp, *Generalized Functionals in Gaussian Spaces: The Characterization Theorem Revisited*, J. Funct. Anal., 1996, 141:301–318.
- [20] A. V. Skorokhod, *On A Generalization of The Stochastic Integral*, Vol. 20, Theory Probab. Appl., 1975, 219–233.

SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY PARK, NOTTINGHAM, NG7 2RD, UK,
E-MAIL: VIACHESLAV.BELAVKIN@NOTTINGHAM.AC.UK

SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY PARK, NOTTINGHAM, NG7 2RD, UK,
E-MAIL: PMXMB1@NOTTINGHAM.AC.UK